

## **GUARDIANS OF HEALTH: ANDROGRAPHIS PANICULATA'S ANTIOXIDANT POTENTIAL EXPLORED THROUGH PHYTOCHEMICAL ANALYSIS AND IN VITRO ASSESSMENT**

**Dr. Sneha R. Kumar and Dr. Priya L. Menon**

Department of Veterinary Pathology, Veterinary College, Bidar.

Department of Veterinary Pathology, Veterinary College, Bengaluru. Vice-chancellor, KVAFSU, Bidar. Director, IAH & VB, KVAFSU, Bengaluru.

### **Abstract**

The balance between free radicals and antioxidants in the body is critical for maintaining overall health. To help address oxidative stress, researchers have been exploring natural, non-toxic compounds with antioxidant potential, such as those found in traditional Indian medicine. One such plant, *Andrographis paniculata*, is the focus of this study, which seeks to analyze its phytochemical composition and assess its in vitro antioxidant capabilities. Results indicate that the ethanolic extract of *Andrographis paniculata* contains a variety of beneficial compounds, including alkaloids, terpenoids, phenolic compounds, tannins, and steroids. Additionally, the extract demonstrated free radical scavenging activity comparable to ascorbic acid (Vitamin C), a well-known antioxidant. These findings suggest that *Andrographis paniculata* holds significant potential as a safe, effective source of antioxidants.

**Keywords:** *Andrographis paniculata*, oxidative stress, phytochemicals, free radical scavenging activity.

### **INTRODUCTION**

The increasing awareness about the role of oxidative stress in various diseases and the importance of antioxidants in preventing these diseases has led to an intense search for natural sources of antioxidants. Oxidative stress, caused by an imbalance between the production of reactive oxygen species (ROS) and the antioxidant defense system, is involved in the pathogenesis of several chronic diseases, including cancer, cardiovascular diseases, diabetes, and neurodegenerative disorders (Valko et al., 2007). Antioxidants function by scavenging free radicals, preventing the formation of ROS, and repairing the damage caused by oxidative stress (Pisoschi et al., 2016). Considering the potential side effects of synthetic antioxidants, there is a growing interest in discovering natural antioxidants from plant sources (Gülçin, 2012). *Andrographis paniculata* (*A. paniculata*), a traditional medicinal plant belonging to the family Acanthaceae, has been used for centuries in Ayurvedic and traditional Chinese medicine for the treatment of various ailments, including fever, respiratory infections, and gastrointestinal disorders (Akbar, 2011). The main bioactive constituent of *A. paniculata* is andrographolide, a diterpenoid lactone, which has been reported to possess significant antioxidant, anti-inflammatory, and anticancer activities (Chua et al., 2017). Apart from andrographolide, several other phytoconstituents, including flavonoids, phenolic acids, and terpenoids, have been identified in *A. paniculata*, which may contribute to its antioxidant potential (Singh et al., 2017). Despite the numerous pharmacological activities reported for *A. paniculata*, there is limited scientific evidence available on its antioxidant potential and the phytochemical profile responsible for this activity. Therefore, the main objective of this study is to explore the antioxidant potential of *A. paniculata* by conducting a comprehensive phytochemical analysis and assessing its in vitro antioxidant activity using various assays. Several methods have been employed to investigate the antioxidant activity of plant extracts, including 2,2-diphenyl-1-picrylhydrazyl

(DPPH) radical scavenging assay, 2,2'-azinobis(3-ethylbenzothiazoline-6-sulphonic acid) (ABTS) radical cation decolorization assay, ferric reducing antioxidant power (FRAP) assay, and oxygen radical absorbance capacity (ORAC) assay (Pisoschi et al., 2016). These assays differ in their mechanisms, sensitivity, and specificity, and it is recommended to use multiple assays to obtain a comprehensive evaluation of the antioxidant potential of a plant extract (Prior et al., 2005). Phytochemical analysis of plant extracts is crucial for understanding the chemical constituents responsible for their biological activities. Various analytical techniques, such as high-performance liquid chromatography (HPLC), gas chromatography-mass spectrometry (GC-MS), and liquid chromatography-mass spectrometry (LC-MS), have been employed for the identification and quantification of phytochemicals in plant extracts (Sultana et al., 2009). In this study, a combination of these analytical techniques will be used to comprehensively profile the phytochemical composition of *A. paniculata*.

In conclusion, this study aims to provide scientific evidence for the antioxidant potential of *A. paniculata* and identify the phytochemical constituents responsible for this activity. The findings from this study may contribute to the development of novel antioxidant agents derived from *A. paniculata* and enhance our understanding of its potential health benefits in the prevention and management of oxidative stress-related diseases.

## **2. MATERIALS AND METHODS**

**2.1 Plant extract and Chemicals:** Dried whole plant ethanolic extract powder of *Andrographis paniculata* was obtained from Himalaya herbal pvt Ltd. Bangalore, India and it was preserved in a sealed vial at 4 °C until tested and analyzed. All other chemicals and reagents used for the study were procured from local sources and were of analytical grade.

**2.2 Preliminary phytochemical tests for the *Andrographis paniculata* extract:** A qualitative phytochemical screening of the ethanolic extract of the *Andrographis paniculata* to detect the presence of essential phytoconstituents, such as alkaloid, tannin, saponin, flavonoid, anthraquinone glycoside, steroids/terpenes, glycosides, proteins, carbohydrates and phenol, was carried out using standard biochemical procedures.

### **2.2.1 Test for Carbohydrates: Benedict test**

2ml of the extract sample mixed with 2ml of the Benedict's reagent and heated in a boiling water bath for 10 minutes. The change in colour to yellow, green to red indicates the presence of reducing sugars.

### **2.2.2 Test for proteins: Biuret test**

3 ml of the extract sample mixed with 1ml of 4% w/v sodium hydroxide and 1ml 1% copper sulphate. The change in colour of the solution to violet or pink indicates the presence of the proteins.

### **2.2.3 Test for alkaloids**

**Wagner's test:** To a few ml of filtrate, few drops of Wagner's reagent was added on the side of the test tube. A reddish – brown precipitate confirmed the test as positive.

**Dragendroff's test:** Dragendroff's reagent was added in a test tube containing 2 to 3 ml of extract filtrate. A prominent yellow precipitate indicated the test as positive

### **2.2.4 Test for Phenols**

**Ferric chloride test:** Two to three drops of 1% Ferric chloride solution was added into 1 mL of extract sample. Phenolic compounds produce a deep violet color or black precipitate with ferric ions.

### **2.2.5 Test for flavonoids: Alkaline reagent test**

2ml of the plant extract was treated with few drops of 20% sodium hydroxide solution. Formation of intense yellow colour which becomes colourless on addition of dilute hydrochloric acid, indicate the presence of flavonoids

### **2.2.6 Test for terpinoids**

**Salkowkis Test:** The Sample was separately shaken with chloroform (2 mL) followed by the addition of concentrated sulphuric acid (2 mL) along the side of the test tube, a reddish-brown coloration of the interface indicates the presence of terpenoid.

#### **2.2.7 Test for tannins**

The sample was stirred with distilled water (10 mL) and then filtered. A few drops of 5% ferric chloride were then added. Black or blue-green coloration or precipitate was taken as positive result for the presence of tannins.

#### **2.2.8 Test for Saponins**

The extract sample (2gm) was shaken vigorously with distilled water (20 mL) in a test tube. The formation of frothing indicates the presence of saponins.

#### **2.2.9 Test for steroids: Libermann Burchard test**

1ml of the plant extract was treated with few drops of chloroform, acetic anhydride and concentric sulphuric acid and observed the formation dark pink or red colour indicates the positive for steroids.

#### **2.2.10 Test for Anthraquinone glycoside**

To the extract solution (1 mL), 5% sulphuric acid (1 mL) was added. The mixture was boiled in a water bath and then filtered. Filtrate was then shaken with equal volume of chloroform and kept to stand for 5 min. Then lower layer of chloroform was shaken with half of its volume with dilute ammonia. The formation of rose pink to red color of the ammonical layer gives indication of anthraquinone glycosides.

### **2.3 In vitro antioxidant activity**

#### **2.3.1 Determination of DPPH scavenging activity**

The antioxidant activity based on scavenging of stable DPPH free radical, was determined as described previously (Mensor *et al.*, 2001). An aliquot of 0.5 ml of each fraction doses (20, 40, 60, 80, 100 and 120 µg/ml) test solution in methanol was mixed with each 2.5 ml of 0.5 mM methanol solution of DPPH. The mixture was shaken well and incubated for 10 min in the dark at room temperature. The absorbance was measured at 517 nm using UV spectrophotometer. Ascorbic acid was used as a positive control and it was mixed in each similar concentration and doses in DPPH as a test solution. All tests and analyses were run in triplicates and the results obtained were averaged. DPPH free radical scavenging ability (%) was calculated by using the formula.

Absorbance of control- Absorbance of sample

% inhibition =  $\frac{\text{Absorbance of control} - \text{Absorbance of sample}}{\text{Absorbance of control}} \times 100$

Absorbance of control

#### **2.3.2 Determination of phosphor–molybdenum scavenging activity**

The antioxidant activity of the extract was determined by the phosphor–molybdenum method as described previously (Prieto *et al.*, 1999). The extract (0.3 ml) of each fractions dose (20, 40, 60, 80, 100 and 120 µg/ml) test solution in methanol of was mixed with 3 ml of reagent solution (0.6 M sulphuric acid, 28 mM sodium phosphate and 4 mM ammonium molybdate). The reaction mixture was incubated at 95°C, for 90 min and cooled to room temperature. Finally, absorbance was measured at 695 nm using a spectrophotometer against blank. Methanol (0.3 ml) in place of extract was used as the blank. Ascorbic acid was used as a positive control and it was mixed in each similar concentration and doses as a test solution. All tests and analyses were run in triplicates and the results obtained were averaged. Phosphor– molybdenum free radical scavenging ability (%) was calculated by using the formula.

Absorbance of control- Absorbance of sample

% inhibition =  $\frac{\text{Absorbance of control} - \text{Absorbance of sample}}{\text{Absorbance of control}} \times 100$

Absorbance of control

### 3. RESULTS AND DISCUSSION

#### 3.1 Phytochemical analysis

The results of our study showed that, ethanolic extract of the *Andrographis paniculata* was found to possess various phytochemicals or polyphenols such as flavonoids, tannins, phenolics, saponins, steroids, carbohydrates, proteins, terpenoids and alkaloids (Table. 1). This result is in agreement with the study carried out by Bajpai *et al.* (2014); Dwivedi *et al.*, (2015); Adegboyega and Oyewole (2015); Sitara *et al.*, (2016) and Rajalakshmi and Cathrine (2016) who reported the presence of carbohydrates, proteins, alkaloids, flavonoids, terpenoids, phenolics, tannins, steroids and saponins. Herbal plants have gained powerful attention due to its effective role in chemo-therapeutic agents. Their prolific effects are mainly due to their phytoconstituents (Kulyal *et al.*, 2010). The various phytochemical compounds detected are known to have beneficial importance in the medical science (Okeke *et al.*, 2001). In recent years, several researchers have reported that phytochemicals including alkaloids, glycosides, terpenoids, saponin, phenols and steroids have enormous antioxidant and free radical scavenging activities (Farhan *et al.*, 2012).

**Table-1: Phytochemical constituents of ethanolic extract of *Andrographis paniculata***

S.No	Phytochemical tests	Results
1	Test for carbohydrates	+ve
2	Test for proteins	+ve
3	Test for alkaloids a. Wagner test b. Drgendroff test	+ve +ve
4	Test for phenolic compounds	+ve
5	Flavonoids	+ve
6	Terpenoids	+ve
7	Tannins	+ve
8	Saponins	+ve
9	Steroids	+ve
10	Anthroquinone glycoside	-ve

#### 3.2 Invitro antioxidant activity

##### 3.2.1 DPPH scavenging activity


DPPH is one of the free radical widely used for testing preliminary radical scavenging activity of a plant extract. In the present study ethanolic extract of AP showed potential free radical scavenging activity in dose dependent manner and statistically significant ( $p < 0.05$ ) along with standard vit-C (Table-2, Figure 1). The scavenging effect of the ethanolic extract of AP and standard vit-C at the concentration of 100  $\mu\text{g/mL}$  were  $69.52 \pm 0.52\%$  and  $80.53 \pm 0.13\%$  respectively.

**Table-2: DPPH Radical scavenging activity (%)**

Concentration (µg/mL)	Extract	Vit-C
20	21.13±0.50 <sup>ax</sup>	38.18±0.47 <sup>ay</sup>
40	31.81±1.56 <sup>bx</sup>	43.06±0.31 <sup>by</sup>
60	40.56±0.23 <sup>cx</sup>	57.55±0.62 <sup>cy</sup>
80	56.69±0.25 <sup>dx</sup>	65.47±0.26 <sup>dy</sup>
100	69.52±0.52 <sup>ex</sup>	80.53±0.13 <sup>ey</sup>
120	80.17±0.42 <sup>fx</sup>	91.84±0.40 <sup>fy</sup>

Values with different superscripts in a row and column vary significantly at  $p < 0.05$

**Figure-1: DPPH Radical scavenging activity (%)**



The DPPH radical is a stable free radical, which has been widely used as sensitive and rapid tool to estimate free radical scavenging activity of both hydrophilic and lipophilic antioxidants (Archana *et al.*, 2005). A freshly prepared DPPH solution exhibits a deep purple colour generally fades/disappears when an antioxidant present in the medium. Thus, antioxidant molecule can quench DPPH free radicals (by providing hydrogen atom or by electron transfer, conceivably via a free radical attack on the DPPH molecule) and convert them to a colourless product (2,2-diphenyl-1-picrylhydrazyl, or a substituted analogous hydrazine) resulting in a decreasing absorbance at 518 nm (Yamaguchi *et al.*, 2002). In the present study, the scavenging effects of the extract and standard vit-C is in increasing trend with increasing concentration of plant extract and Vit-C. The results were concentration dependent and similar findings on DPPH radical scavenging activities of plant extracts have been previously observed (Farhan *et al.* 2012; Amari *et al.* 2014).

### 3.2. 2. Phosphor-molybdenum scavenging activity


The invitro evaluation of the AP extract for its antioxidant potential revealed that the scavenging potential of the extract and standard vit-C for Phosphomolybdenum radical is in increasing trend with increasing concentration of the extract and standard vit-C (Table 3 Figure 2) and statistically significant ( $p < 0.05$ ). The scavenging effect of the ethanolic extract of AP and standard vit-C at the concentration of 100 µg/mL were  $63.60 \pm 0.64$  % and  $67.13 \pm 0.54$  % respectively.

**Table-3: Phosphomolybdenum total antioxidant activity (%)**

Concentration(µg/mL)	Extract	Vit-C
20	7.46±0.35 <sup>ax</sup>	11.40±0.41 <sup>ay</sup>

40	22.40±0.23 <sup>bx</sup>	34.86±0.50 <sup>by</sup>
60	40.26±0.29 <sup>cx</sup>	49.46±0.30 <sup>cy</sup>
80	53.60±0.87 <sup>dx</sup>	61.13±0.35 <sup>dy</sup>
100	63.60±0.64 <sup>ex</sup>	67.13±0.54 <sup>ey</sup>
120	70.93±1.04 <sup>fx</sup>	75.16±0.24 <sup>fy</sup>

Values with different superscripts in a row and column vary significantly at  $p < 0.05$  **Figure-2: Phosphomolybdenum total antioxidant activity (%)**



PM assay is based on the reduction of the phosphate-Mo (VI) to phosphate Mo (V) by the sample and subsequent formation of a bluish green coloured phosphate/Mo complex at acidic pH. Phosphomolybdenum method is routinely applied in the laboratory to evaluate the total antioxidant capacity of plant extracts (Prieto *et al.*, 1999). In the present study, the total antioxidant activity of the extract is in increasing trend with increasing concentration of the extract and standard vit-C. Similar findings on Phosphomolybdenum method for total antioxidant activities of various plant extracts have been previously observed (Yadav, 2015; Phatak and Hendre, 2014; Sangeeta and Venkatlakshmi, 2017).

#### 4. CONCLUSION

On the basis of the results obtained in this study, it is concluded that, the ethanolic extract of *Andrographis paniculata* which contain alkaloids, flavonoids, terpenoids, tannins, saponin, phenols and steroids as essential phytochemicals, exhibited significant antioxidant and free radical scavenging activities. These findings indicate that *Andrographis paniculata*, can be useful therapeutic agent for prevention of oxidative stress.

#### 5. REFERENCES

- Adegboyega, A. and Oyewole, B., 2015. Phytochemical screening and antimicrobial activities of leaf extracts of *Andrographis paniculata*. *Int. J. Sci. Res.*, 4(6): 6-14.
- Amari, N.O., Bouzouina, M., Berkani, A. and Lotmani, B., 2014. Phytochemical screening and antioxidant capacity of the aerial parts of *Thymelaea hirsuta* L. *Asian Pac. J. Trop. Dis.*, 4:104-109.
- Archana, B., Dasgupta, N. and De, B. 2005. In vitro study of antioxidant activity of *Syzygium cumini* fruit. *Food Chem.* 90: 727-733.

- Bajpai, V.K., Agrawal, P. and Park, Y.H., 2014. Phytochemicals, Antioxidant and AntiLipid Peroxidation Activities of Ethanolic Extract of a Medicinal Plant, *Andrographis paniculata*. *J. Food Biochem.*, 38(6): 584-591.
- Chen, H. W., Huang, C. S., Liu, P. F., Li, C. C., Chen, C. T., Liu, C. T., ... & Lii, C. K. 2013. *Andrographis paniculata* extract and andrographolide modulate the hepatic drug metabolism system and plasma tolbutamide concentrations in rats. *Evidence-Based Comp. Altern. Med.*, 2013.
- Dwivedi, D., Thanwar, M and Gharia, A., K. 2015. A phytochemical investigation on *Andrographis paniculata*. *J. Chem. Pharm. Res.*, 7(10): 822-827.
- Farhan, H., Malli, F., Rammal, H., Hijazi, A., Bassal, A., Ajouz, N. and Badran, B. 2012. Phytochemical screening and antioxidant activity of Lebanese *Eryngium creticum* L. *Asian Pac. J. Trop. Biomed.* 2: S1217–S1220.
- Hossain, M. D., Zannat Urbi, Abubakar Sule, and HAFIZUR Rahman. 2014. *Andrographis paniculata* (Burm. f.) Wall. ex Nees: A Review of Ethnobotany, Phytochemistry, and Pharmacology. *Sci. World J.*, **2014**: 1-28 <https://doi.org/10.1155/2014/274905>
- Kulyal., P., Tiwari, U., Shukla, A., and Gaur, A. 2010. Chemical constituents isolated from *Andrographis paniculata*. *Indian J. Chem.*, 46B. 356.
- Mensor, L.L., Menezes, F.S., Leitao, G.G., Reis, A.S., dos Santos, T.C., Coube, C.S., and Leitao, S.G., 2001. Screening of Brazilian plant extracts for antioxidant activity by the use of DPPH free radical method. *Phytother. Res.* 15:127–130.
- Okeke, M.I., Iroegbu C.U., Eze, E.N., Okoli, A.S., and Esimone, C.O. 2001. Evaluation of the Extracts of the Roots of *Landolphia Owerrience* for Anti-bacterial activity, *J. Ethanopharmacol*, 78: 119-127.
- Patel, A., Patel, A., Patel, A. and Patel, N.M. 2010. Determination of polyphenols and free radical scavenging activity of *Tephrosia purpurea* linn leaves (Leguminosae). *Pharmacogn. Res.*, 2, 152–158.
- Phatak R.S and Hendre A.S., 2014. Total antioxidant capacity (TAC) of fresh leaves of *Kalanchoe pinnata*. *J. Pharmacogn. Phytochem.* 2 (5): 32-35.
- Prieto, P., Pineda, M., and Aguilar, M., 1999. Spectrophotometric quantization of antioxidant capacity through the formation of a phosphor–molybdenum complex: specific application to the determination of vitamin E. *Anal. Biochem.* 269: 337–341.
- Rajalakshmi, V. and Cathrine, L., 2016. Phytochemical screening and antimicrobial activity of ethanolic extract of *Andrographis paniculata*. *J. Pharmacogn. Phytochem.*, 5(2): 175.
- Sangeeta, M and Venkatalakshmi, P. 2017. In vitro antioxidant activity of the Aqueous extract of *Andrographis paniculata* and *Carica papaya* leaves. *World J Pharm Sci.*, 6(5): 163116433.

- Sithara, N.V., Komathi, S., Rajalakshmi, G., Queen, J. and Bharathi, D., 2016. Phytochemical analysis of *Andrographis Paniculata* using different solvents. *Phytochemical analysis*, 4(8): 7-10.
- Trivedi N, P. and Rawal, U. M. 2001. Hepatoprotective and antioxidant property of *Andrographis paniculata* (Nees) in BHC induced liver damage in mice. *Indian J. Exp. Biol.*, **39**: 4 1-46.
- Akbar, S. (2011). *Andrographis paniculata*: A review of pharmacological activities and clinical effects. *Alternative Medicine Review*, 16(1), 66-77.
- Chua, L. S., Yap, K. C., & Jaganath, I. B. (2017). An update on the antioxidant capacities and total phenolic contents of *Andrographis paniculata*. *International Journal of Food Properties*, 20(S3), S1459-S1474.
- Gülçin, İ. (2012). Antioxidant activity of food constituents: An overview. *Archives of Toxicology*, 86(3), 345-391.
- Pisoschi, A. M., Pop, A., Cimpeanu, C., & Predoi, G. (2016). Antioxidant capacity determination in plants and plant-derived products: A review. *Oxidative Medicine and Cellular Longevity*, 2016, 9130976.
- Prior, R. L., Wu, X., & Schaich, K. (2005). Standardized methods for the determination of antioxidant capacity and phenolics in foods and dietary supplements. *Journal of Agricultural and Food Chemistry*, 53(10), 4290-4302.
- Singh, R., Mishra, R. K., & Singh, S. (2017). Chemical composition and bioactivity of *Andrographis paniculata* (Burm. f.) Nees. *Indian Journal of Pharmaceutical Sciences*, 79(2), 286-292
- Sultana, B., Anwar, F., & Przybylski, R. (2009). Antioxidant activity of phenolic components present in barks of *Azadirachta indica*, *Terminalia arjuna*, *Acacia nilotica*, and *Eugenia jambolana* Lam. *Trees. Food Chemistry*, 104(3), 1106-1114
- Valko, M., Leibfritz, D., Moncol, J., Cronin, M. T., Mazur, M., & Telser, J. (2007). Free radicals and antioxidants in normal physiological functions and human disease. *The International Journal of Biochemistry & Cell Biology*, 39(1), 44-84.
- Wasman, S.Q., Mahmood, A.A., Chua, L.S., Alshawsh, M.A. and Hamdan, S., 2011. Antioxidant and gastroprotective activities of *Andrographis paniculata* (Hempedu Bumi) in Sprague Dawley rats. *Indian J. Exp. Biol.*, **40** (10): 767-772.
- Yadav Yogesh C. 2015. Hepatoprotective effect of *Ficus religiosa* latex on cisplatin induced liver injury in Wistar rats. *Revista Brasileira de Farmacognosia.*, 25:278–283. [21] Yamaguchi, T., Takamura, H., Matoba, T. and Terao, J., 2002. HPLC method for evaluation of the free radical scavenging activity of foods by using 2, 2-diphenyl-1-picryl hydrazyl. *Bio Sci Biotechnol Bio Chem.*, 62: 1201-1204.