

RÍO SOTO LA MARINA'S HIDDEN GEM: FUNDULUS HERMINIAMATILDAE, A NOVEL KILLIFISH SPECIES DISCOVERED IN THE UPPER BASIN OF NUEVO LEÓN, MEXICO

Prof. Mariana L. González and Dr. Alejandro M. García

Laboratorio de Ictiología, Facultad de Ciencias Biológicas, Universidad Autónoma de Nuevo León,
MEXICO

Abstract

Fundulus herminiamatildae sp. nov. Is endemic from the Marmolejo stream, a head water tributary of Río Soto La Marina basin, in the municipality of Aramberri, Nuevo León, México. Geologically, it is located in the Northeastern province of México, specifically in the Sierra Madre Oriental subprovince. The water temperatura is a determinant and important factor for the species differentiation. Its closest relative is *F. philpisteri*. *Fundulus herminiamatildae* is distinguished from other killifishes by a large number of conspicuous and simple lateral bars, body with high profile, and the following proportions in cephalic length: snout (2.52-9, mean 2.7) eye diameter (4.1-4.9, mean 4.5), and Preorbital length (2.7-3.1, mean 2.9).

Keywords: *Fundulus herminiamatildae*, Mexico, Marmolejo stream, Río Soto La Marina, New species.

Acknowledgments

The authors thank Dr. Anthony Echelle and Dr. Gorgonio Ruiz-Campos for the corrections made to the manuscript. Our thanks also go to the collectors of the specimens: A. J. Contreras-Balderas, J. A. Contreras-Lozano, J. E. Lozano-Vilano, and T. Rodriguez; to Biol. Hidalgo Rodríguez Vela for designing the map; to José Luis Gibaja for photos; to technician A. Ovalle Segovia for his assistance in the laboratory; and to T. Guadarrama for helping with the bibliographic material; and Universidad Autónoma de Nuevo León (PAICYT, contract CN 040-09). The authors are also thank ful to every one who has contributed their part in carrying out this research.

The genus *Fundulus* Lacépède, 1803 belongs to the American cyprinodontiform family, Fundulidae (Parenti, 1981), of Gondwanan origin (Nelson, 2006). This genus includes euryhaline and eurythermic species in habiting desertic and semi-desertic areas, springs, rivers, and lagoons of North America, including coastal areas in México. It is composed of approximately 40 species (Miller, 1955; Mayden et al. 1992; Nelson, 1994; García-Ramírez et al. 2006) among which eight are found in Mexico: *Fundulus grandis* Baird & Girard, 1853 in States of Tamaulipas and Veracruz; *F. grandissimus* Hubbs, 1936 in North of Yucat an and Quintana Roo, *F. persimilis* Miller, 1955 Northern of Yucatan; *F. lima* Vaillant 1894; *F. parvipinnis* Girard, 1854 Baja California peninsula; *F. philpisteri* García-Ramírez et al., 2006 in Nuevo León; *F. similis* (Baird & Girard, 1853) in Tamaulipas and *F. zebrinus* Jordan & Gilbert, 1883 an exotic species in Mexico, found in Chihuahua. Currently, *F. grandissimus* and *F. persimilis*, are respectively considered as vulnerable and endangered species in the IUCN Red List of Threatened species (Collette, B, et al. 2015; Jelks, H., et al. 2015) and are under a special protection in México by NOM-059-ECOL-2001 (Norma Official Mexican). In addition, *F. philpisteri* presents amore alarming status, it is classified as a critically endangered species (Maíz-Tome, L. 2019) due to its fragile habitat and specialization. In México García-Ramírez (1997) reported that the population of Arroyo Marmolejo was different from *Fundulus grandis*.

The aim of this paper is to provide a morphological description of a new species (*F. herminiamatildae*) on the basis of specimens from a head water stream (ArroyoMarmolejo) of the Río Soto La Marina basin, in the municipality of Aramberri, Nuevo León, México.

Additionally, we compared morphological character soft his new species with those of termal spring (*F. philpisteri*) and estuarine/fresh water environments (*F. grandis*) in order to determine diagnostic and description characteristics.

2. Methods

The specimens examined of the genus *Fundulus* for this study were collected along coastal environments, comprising the Laguna Madre Tamaulipas system, including its associated streams, and in headwater streams of the San Fernando and Soto La Marina river basins, between 1966-2004 (Fig. 1). The examined material was housed at the Fish Collection of the Facultad de Ciencias Biológicas, Universidad Autónoma de Nuevo León (UANL) and several paratypes were deposited indifferent institutions: Museum of Zoology University of Michigan (UMMZ-248753), Tulane Museum of Natural History (TUHC-204176), University of Texas at Austin (TNHC-39831), Escuela Nacional De Ciencias Biológicas, Instituto Politécnico Nacional (IPN-ENCB-P5592), United States National Museum (USNM-391644).

Abbreviations for institutions and collections cited follow Leviton et al. (1985). The examined specimens were catalogued with the acronymous, number of catalog, number of specimens, standard length as minimum and maximum in millimeters, locality, basin, state, country, collector(s), and collecting dates were added.

Thirty-five body measures and 16 meristic characters of the standardized method of Hubbs and Lagler (1958) were considered for the morphometric analysis of the specimens (Fig.4). All the measurements were made on the left side of each specimen using a digital caliper (precision, 0.01 mm). Proportions are expressed in base of the TSL= times in standard length; TCL = times in cephalic length. Count sof the cephalic pores follow Gosline(1949) were considered.

For a multivariate approach a canonical discriminant function was realised using SPSS (version10.0) was performed. To carry out this analysis, we examined 20 specimens of *Fundulus grandis* (Laguna Madre, Tamaulipas), 18 specimens of *F. philpisteri* (Baño de San Ignacio, Linares, Nuevo León) and 15 from specimens of *F. herminiamatildae* (new species of Marmolejo stream, Aramberri, Nuevo León).

2.1. Comparative material. *Fundulus philpisteri*. - Baño de San Ignacio springs, Linares, Nuevo León, México, Río San Fernando: UANL 15031 (26:28-83.7mm); Coll.: S. Contreras-Balderas and grupo de la Universidad Autónoma de Tamaulipas; 7 Sep. 2001; UANL 9031 (1: 63.4 mm); Coll.: S. Contreras-Balderas and A. Contreras-Arquieta; 6 Aug. 1988; UANL 11146 (5:31.6-37.3 mm); Coll.: S. Contreras-Balderas, M. L. Lozano-Vilano, M. E. García-Ramírez, and A. J. Contreras-Balderas, 12 oct. 1992; UANL-16147 (1:

56.1mm); Coll.: M. L. Lozano-Vilano, A. J. Contreras-Balderas, Jorge A. Contreras-Lozano and J. E. LozanoVilano, 23 jul. 2004.

Fundulus grandis. - Río Bravo: UANL 2138 (34:35.2-71.0 mm) Presa Falcón cerca de Nueva Cd. Guerrero, Tamaulipas, México; Coll.: S. Contreras-B. and Gpo. de Fac. de Biología, 29 Jul. 1975; UANL 1523 (4:31.3-48.4mm) Presa Marte R. Gómez, 6 km NNW of Comales; Coll.: S. Contreras-B. and Gpo. FCB, 24 Sep. 1973; UANL 6154 (25:50.6-61.3 mm) Río Bravo en Miguel Alemán; Coll.: S. Contreras-B. and Gpo. FCB, 30 Jul.1982; UANL 6097 (1: 31.6 mm) Río Bravo en Matamoros; Coll. S. Contreras-B. and Gpo.

FCB, 28 Jul. 1982; Coll.: S. Contreras-B., A. Gómez-S., and Gpo. Biol. de Campo, 8 Sep. 1990; UANL 11454 (13:66.5-81.4 mm) Río San Juan bajo puente Carr. México 2km W of Camargo, México; Coll.: A. J.

Contreras-B., P. J. Rodríguez-B. and J. A. Bernal, 1 Apr. 1977; UANL 2126 (1: 35.0 mm) Boca del Río Bravo;

Coll.: S. Contreras-B. and Gpo. FCB., 28 Jul. 1975; UANL 4181 (1: 47) Río Álamo en la desembocadura al Río Bravo, México; Coll.: G. Ruíz-C., H. Cirilo-S., H. Torres-M. and D. J. Castillo-R., 18 Dec. 1981; UANL 6128 (3:21.6-47.0 mm) Río Bravo en Parque Anzaldúas, México; Coll.: S. Contreras-B., S. Contreras-A., A.

Contreras-A. and A. Contreras-A., 29 Jul. 1982. Laguna Madre; UANL 7936 (7: 57.4-83.8 mm) Punta de la

Media Luna (No San Luis Potosí), México; Coll.: A. Gómez-S, C. Rosario-R., and J. Rabago-C, 24 May 1986; UANL 10466 (18:37.2-50.8 mm) Laguna Almagre, México; and UANL 5320 (9:61.7-95.5 mm) Río San Juan en el Salto, 6 km NE de los Aldabas, Nuevo León, México; Coll.: C.M. Villarreal-T., M. E. Obregón-M. and J. F. Lozano-M., 3 May 1982.

3. Results

FUNDULUS HERMINIAMATILDAE SP.NOV.

(Table I, Fig. 3) Material examined

3.1. Holotype. - UANL-18803, adult male, 88.7 mm SL; Marmolejo stream, Aramberri, Nuevo León, México, Soto La Marina River basin (24° 09' 38.6" N 99° 45' 15.7" W); Coll.; S. Contreras-Balderas, M. L. Lozano-Vilano, M. E. García-Ramírez, A. J. Contreras-Balderas and T. Rodríguez; 28 Apr. 1992.

3.1.1. Paratypes. - UANL-13613 (36: 106.4-30.4 mm SL), same data as holotype, UANL-16160 (29: 87.4-32.0 mm SL) stream at SW of El Porvenir, Nuevo León, México; Coll.: M.L. Lozano-Vilano, A. J. Contreras-Balderas, J. A. Contreras-Lozano, and J. E. Lozano-Vilano; 25 Jul. 2004; UANL-16161 (22: 55.140.1 mm SL) small pond at the SW of El Porvenir; Coll.: M. L. Lozano-Vilano, A. J. Contreras Balderas, J.A.

Contreras-Lozano and J.E. Lozano-Vilano; 25 Jul. 2004; UANL 6448 (7: 38-67 mm LP) Río Blanco at 7 Km

N of Aramberri; Coll.: Salvador Contreras-B., H. Obregón-B. and A. Contreras-A., 20 Nov.1985; UANL 6447(35:37-86) RíoBlanco 4km from Aramberri; Coll.: S. Contreras-B., H. Obregón-B. and A. Contreras-A., 20 Nov. 1985; UANL 1149 (16:36-83) Río Aramberri, 3km NE of Aramberri; Coll.: F. Jiménez-G., A.

Rodríguez, and R. Banda, 7 Aug. 1968; UANL 1133 (145:29-86) 5 km NE of Aramberri; Coll.: S. ContrerasB., D. Molina, and J. A. Martínez, 24 Apr. 1966; UANL 13617 (21:29-79) Arroyo 4 km E of Aramberri; Coll.: S. Contreras-B and Grupo FCB, 13 Dec. 1987; UANL 15696 (41: 72.6-26.9) Arroyo Aramberri; Coll.: M.L. Lozano-V., A. J. Contreras-B., J. A. Contreras-L., and J. E. Lozano-V., 25 Jul. 2004; UANL 16160 (29: 89.1-30.4) Arroyo al SW del Porvenir; Coll.: M. L. Lozano-V., A. J. Contreras-B., J. A. Contreras-L. and J. E. Lozano-V., 25 Jul. 2004; and UANL 16161 (22: 57.3-33.7) Charco al SW del Porvenir; Coll.: M. L. LozanoV., A. J. Contreras-B, J. A. Contreras-L. and J. E. Lozano- V., 25 Jul. 2004.

4. Discussion

4.1. Diagnosis. *Fundulus herminiamatildae* is distinguished from the other species in northeastern Mexico, using the following combination of characters: 12-17 conspicuous and simple lateral bars, subangular head, thickened body, and slightly convex predorsal profile. Proportions in SL of the new species (Table II) and its comparison with other relatives is as follows: head length of 2.9-3.5 (3.2)

(mean is shown in parenthesis), versus 3.0-3.3 (3.2) in *F. grandis* and 2.8-3.2 (3.0) in *F. philpisteri*; anal fin origin-hypural length of 2.6-3.0 (2.8) versus 2.6-2.8 (2.6) in *F. grandis* and 2.7-3.3 (3.0) in *F. philpisteri*; caudal peduncle short of 3.9-4.2 (4.1) versus 3.4-4.1 (3.8) in *F. grandis* and 3.9-4.7 (4.2) in *F. philpisteri*; snout-pelvic origin length of 1.7-2.0 (1.8) versus 1.9-2.1 (2.0) in *F. grandis* and 1.8-2.2 (1.9) in *F. philpisteri*; dorsal-anal origin length of 3.7-4.3 (3.9) versus 3.7-4.6 (4.1) in *F. grandis* and 3.9-4.9 (4.3) *F. philpisteri*; least depth short of 5.4-6.9 (6.1), versus 5.7- 6.4 (6.2) in *F. grandis* and 6.0-7.1 (6.6) in *F. philpisteri*.

Proportions in head length (HL, Table II) of the new species are pelvic-pectoral origin length of 3.5-4.5 (4.1) versus 4.8-5.3 (5.0) in *F. grandis* and 3.6-5.0 (4.4) in *F. philpisteri*; the organism is also characterized by big snout of 2.5-2.9 (2.7) versus 3.1-4.0 (3.5) in *F. grandis* and 2.5-3.4 (3.2) in *F. philpisteri*; small eyes of 4.1-4.9 (4.5) versus 3.8-4.4 (4.0). In *F. grandis* and 3.1-4.8 (4.0) in *F. philpisteri*; preorbital width of 2.7-3.1 (2.9) versus 3.2-3.7 (3.4) in *F. grandis* and 3.0-3.8 (3.4) in *F. philpisteri*; anal length depressed to 1.3-1.9(1.5) versus 1.3-1.4(1.3) in *F. grandis* and 1.3-2.1(1.6) in *F. philpisteri*.

The canonical discriminant analysis showed that the three species compared here can be separate completely (Fig. 2). A total of 35 morphometric variables on 15 specimens of *F. herminiamatildae*, 18 of *F. philpisteri*, and 20 of *F. grandis* were compared and analyzed through discriminant analysis. The Wilk's Lambda values changed from 0.001 to 0.056, which was interpreted as firm discrimination among the compared species.

Canonical functions 1 (F1) and 2 (F2) explained 75.2% and 24.8% of variance ($p < 0.000$) respectively.

4. 1. 1. Description. The physiognomy of the Holotype is shown in figure 3 and its morphometric characters are presented in tables I and table II with comparison of more distinctive features in proportional measures in the three species. The characters observed in both the sexes include wide preorbital 2.7-3.1(2.9) times of the head length (the average is shown in parenthesis); small eyes of 4.1-4.9 (4.5) times of the head length, located closer to the snout than the operculum, snout projected upward, dorsal fin inserted slightly in front of the anal origin, reaching three-fourths of the caudal peduncle, flattened occipital region, rounded anal dorsal, pectoral, and pelvic fins.

The following characteristics were observed among males: thickened body, with body depth higher than the females; nuptial tubers above the dorsal and ventral zone; dorsal, anal, pectoral, and caudal fins; long dorsal and anal fins projected three-fourths over the caudal peduncle; and pectoral fins extended until or beyond the anus.

The females possess elongated body; short pelvic fins, not reaching the anus; and dorsal fin extended to the half of the caudal peduncle.

The meristic data are as follows (holotype in parenthesis): fin rays: dorsal 9-11(11); anal 9-11(11); pectorals 15-19 (18); pelvics 5-7 (7); and caudal 18-20(18). Scales: lateral Line 35-38 (37); dorsal to anal 12-15(13); dorsal-pelvic insertion 13-15(13); predorsals 21-24(21); around caudal peduncle 21-27(22); around the body 31-37(35); gill rakers 9-12 (9); and head pores: mandibulars 5(5); cephalic 8-9(9), lacrymals 4(4); and Preoperculars 7(7).

4.1.1. 1. Coloration. Live coloration.- Mature males with body and posterior part of dorsal fin olive with white spots; pectorals, pelvics and anal fins orange, anal with white spots, caudal yellowish or olive with white spots, body side with 13-17 conspicuous dark bars (juveniles with bars more evident).

Mature females with head bluish, body yellowish and ventral region cornsilk, sides with 13-17 conspicuous dark bars; all fins oranges.

4.1.2. Preserved specimens.- Males body well-pigmented in the dorsal región with dark-brown head; orange ventral region) body and dorsal fin, anal and caudal with colorless spots; 13-17 simple and double conspicuous lateral bars (better marked in juveniles than in adults).

Females: melanophores in the anterior part of the body and head.

4.1. 2. 2. Sexual dimorphism. Males have deeper bodies and generally longer fins than females. Nuptial tubercles presents in body and fins. Females as in the other members of the species flock had a membranous bag extended to the third or fourth anal radius.

4.1. 2. 2. 2. Habitat and associates. The Marmolejo área is located 11 km downstream the river of the city hall of Aramberri, Nuevo León, in the physiogeographical subprovince of plains and hills (SPP, 1985). The average altitude of the área is 930 masl. The local fish assemblage of *F. herminiamatildae* sp. nv. Comprises *Xiphophorus xiphidium* and *Cyprinella* sp. The hábitat consist of cold mountain streams with rocky bottoms and riparian vegetation composed by syxamore trees (*Platanus occidentalis*) and mule fat (*Baccharis salicifolia*).

4.1.3. Etymology. The scientific name of *herminiamatildae* is patronymic for Herminia Ramírez and Matilde García, parents of the main author, who were important pillars in the profesional and personal growth of the author, and who have a great respect for nature.

5. 1.3.1 Distribution. *Fundulus herminiamatildae* sp. nov. is endemic to the surrounding área of Aramberri, in the upper Soto La Marina River basin, Nuevo León, México (24°09'38.6"N and 99°45'15.7"W).

References

- Collette, B., Grubbs, D., Pezold, F., Simons, J., Carlson, J., McEachran, J. D., Brenner, J., Tornabane, L.,
- Chakrabarty, P., Roberson, R., Chao, L., Tolan, J., Espinosa-Perez, H., Vega-Cendejas, M., & Jelks, H. (2015). *Fundulus grandissimus*. The IUCN Red List of Threatened Species 2015: e.T191311A1975554. <http://dx.doi.org/10.2305>.
- García-Ramírez M. E., (1997). Revisión Sistemática del Complejo *Fundulus grandis* (Pisces: Fundulidae) del Noreste de México. Tesis de Maestría Inédita, 128 p. Univ. Autónoma de Nuevo León, México.
- García-Ramírez M.E., Contreras-Balderas S., & Lozano-Vilano M.L. (2006). – *Fundulus philpisteri* sp. nv. (Teleostei: Fundulidae) from the río San Fernando Basin, Nuevo León, México. In: M. L. Lozano-Vilano & A. J. Contreras Balderas., (Eds) *Studies of North American Desert Fishes in Honor of E.P. (Phil) Pister Conservationist*. Universidad Autónoma de Nuevo León, Monterrey, México, (pp13-19).
- Gosline, W.A. (1949). The sensory Canals of the head in some Cyprinodont fishes, with particular reference to the genus *Fundulus*. *Occ. Pap. Mus. Zool. Univ. Mich.*, 519:1-17.
- Hubbs, C L., & Karl F. Lagler (1958). *Fishes of the Great Lakes Region*. Cranbrook Inst. Sci. Bull., 26:i-xi, 1-213 (2d. Ed.)

Jelks, H., Espinosa-Perez, H., Vega-Cendejas, M., & Tolan, J. (2015). *Fundulus persimilis* The IUCN Red List of

Threatened Species 2015: e. T169374A1274789.<http://dx.doi.org/10.2305>

fishes (Telostei:Atherinomopha) Bull. Am. Mus. Nat. Hist. 168:341-557.

Secretaría de Programación y Presupuesto. México.