

## **NUTRITIONAL STRATEGIES FOR ENHANCING GROWTH IN WEANED LAMBS: A CONCENTRATE STUDY**

**Dr. Christopher David Walker<sup>1</sup>, Prof. Rachel Nicole Johnson<sup>2</sup> and  
Dr. Marcus Anthony Thomas<sup>3</sup>**

Caroni North Bank Road, Centeno, University of Trinidad and Tobago, ECIAF Campus<sup>1</sup>

University of Trinidad and Tobago, Eastern Caribbean Institute of Agriculture and Forestry, Faculty of  
Biosciences<sup>2</sup>

Agriculture and Food Technology, Caroni North Bank Road, Centeno Arima, Trinidad<sup>3</sup>

---

### **Abstract**

The small ruminant livestock industry in Trinidad and Tobago faces significant challenges in terms of productivity, primarily attributed to poor reproductive strategies, limited genetic diversity, and inadequate veterinary support. However, the most critical factor affecting productivity is the suboptimal feed supply and feeding practices. The available feed resources, primarily natural pasture and crop residues, are characterized by their high fibrous content, substantial lignin levels, and overall low nutritive value. Animals consuming such low-quality feeds tend to utilize their fat reserves to meet nutritional needs, resulting in reduced body fat storage, particularly exacerbated during periods of forage scarcity. Livestock's live weight and nutritional status significantly influence key production efficiency metrics like feed conversion ratio (FCR) and average daily gain (ADG). Dry forages and roughages, with a crude protein (CP) content below 7%, often fail to meet microbial requirements unless supplemented with protein-rich feeds. When utilized in isolation, these poor-quality forages are insufficient even to meet the maintenance needs of livestock. Continued reliance on such low-quality feed sources, including lignified grasses and fibrous crop residues, yields negligible improvements in live weight gain and overall performance, particularly for weaned lambs. Consequently, an urgent need exists to enhance the feeding system for small ruminants, with special emphasis on supplementing grasses with high-energy feeds.

**Keywords:** Small Ruminant Livestock, Feed Supply, Nutritional Status, Livestock Productivity, Feed, Conversion Ratio.

---

### **Introduction**

Poor reproductive strategies, inadequate genetic stock, the lack of veterinary services and inadequate quantity and quality of feed can be listed as the causes for low productivity of small ruminant livestock in Trinidad and Tobago. Among these limiting factors, poor feed supply and feeding system is the most critical as the feed resources in Trinidad are comprised largely of natural pasture and residues of different crops. McDonald *et al.* (2002) indicated that all the straws, and related by-products are extremely fibrous, most of them have high levels of lignin and all are of low nutritive value. It was reported that animals fed low quality feed use their fat body reserve in order to fulfill their nutrient requirement which in turn leads to decreased fat storage in their body and as a result mobilization of body fat reserves will start. This is pronounced during dry periods of forage growth and prolonged underfeeding Kirton *et al.* (1995). Kirton *et al.* (1995) reported that live weight and nutritional status of

animals can influence their production efficiency factors; such as feed conversion ratio (FCR) and average daily gain (ADG). Most dry forages and roughages possess a crude protein (CP) content of less than 7 % which indicates microbial requirement can hardly be satisfied unless supplemented with feeds rich in protein (VanSoest *et al.*, 1994). When fed alone, poor quality forage are unable to provide even the maintenance requirement of livestock (Rege, 1999). Continued feeding of low quality feed stuffs like lignified grass and fibrous crop residues will not result in increases in the rate of live weight gain and enhance performance of weaned lambs. Therefore, it is important to consider the improvement of the feeding system for small ruminants. In this case, the use of supplementation of grass with high energy feeds need due attention.

One possible method of improving the nutritive values of crop residues is through the strategic supplementation with energy and protein rich sources that has the capacity to promote voluntary feed intake, digestibility and subsequent utilization (Nega and Melaku, 2009).

Hence, there is a need to search for alternative feeding methods, which could supplement poor quality roughage feeds to enhance productivity of livestock in the tropics. Some researchers have indicated that lambs supplemented with different levels of concentrate mixture were able to accommodate relatively better lean flesh than lambs fed wheat straw alone. This finding was in line with many other studies, (Asnakew, 2005; Mulu, 2005; Simret, 2005; and Zemichael, 2007). Payne and Wilson (1999) also reported that dressing percentage increases with increasing the proportions of concentrate in the ration. Sandros (1993) reported that grazing lambs supplemented with concentrate had significantly higher slaughter weight, hot carcass weight, and dressing percentage than the non-supplemented lambs. Apparent nutrient digestibility of dry matter (DM) and organic matter (OM) of the basal diet was significantly increased by 14.2% of dry matter (DM) due to supplementation. In line with this result Khanal *et al.* (1999) reported that there was a 13.3% increment in apparent dry matter (DM) digestibility for sheep supplemented with concentrate mixtures.

The demand for fresh lamb meat in the local market has increased over the last decade even with a decline in local production. Local producers have developed a feeding system where poor quality grass is supplemented with heavy amounts of concentrate to fatten lambs. Feeding accounts for 70 – 80 % of production cost. Escalating feed prices coupled with our present economic climate have forced local producers to adjust the utilization of concentrate feeds. This study was therefore designed to investigate an appropriate concentrate inclusion rate that could positively impact the performance parameters of growing lambs in the most cost effective manner.

## **Materials and methods**

### **Experimental animals, feeds and management**

This study was conducted at the Eastern Caribbean Institute of Agriculture and Forestry Farm, located in Caroni North Bank Road Centeno. The trial's method and procedures were accepted by the Ethics Committee of the University of Trinidad and Tobago. A total of 21 female Barbados Black Belly lambs, weighing an average of 15 to 19 kg was used to evaluate the effect of dietary inclusion of a commercial concentrate (Ruminant mix) on growth performance. Lambs were treated against internal parasites before commencement of the experiment. Seven lambs each of a group of 21 were randomly allotted to

each of three dietary treatment groups, that is; Treatment 1: inclusion @ 2 % body weight, Treatment 2: inclusion @ 2.5 % and Treatment 3: inclusion @ 3 % body weight. Animals were subjected to the experimental diets for an adaptation period of 12 days before the commencement of the experiment. The amount of feed offered each time to the lambs were recorded daily. Refusals and wastage were weighed on the following morning. The total weight of the feed offered, adjusted for refusal and wastage, provided an estimate of the average feed consumed by each group per day throughout the study. The animals had free access to water and the volume of water offered each time was recorded daily and the remaining was recorded on the following morning. All of the lambs were fed a basal diet of mixed grass harvested from the farm. Daily water intake, corrected for losses, provided an estimate of the average water intake per day for each group. Animals were weighed on a weekly basis before offering fresh feed. Samples of feed were collected biweekly for proximate analyses (AOAC, 1995).

### **Chemical Analyses**

The DM content was measured by placing 1 g of sample in a porcelain crucible in an oven set at 100°C overnight. The loss in weight was used as a measure of moisture content. Organic matter was determined by igniting 1 g of sample in a muffle furnace at 550 °C and estimated as the loss in weight of sample. Nitrogen content was determined by using the Kjeldahl method (AOAC 1995, method no.990.03) and was converted to crude protein by multiplying percentage N content by 6.25. Ether extract was determined in accordance with the Soxhlet extract method using petroleum ether as the extract agent (AOAC, 1995). Ash content was determined by incinerating the samples in a muffle furnace at 550 °C (AOAC, 1995).

### **Carcass analysis**

At the end of the experiment, 3 animals were selected from each of the treatments and starved overnight, while having access to water. Live weights at slaughter and dressed carcass weights were recorded.

### **Economic efficiency**

This was evaluated as the feed cost for each pound of weight that was gained by the lambs. Hence, efficiency was calculated as the ratio between income (price of weight gain) and the cost of feed consumed (**feed cost ÷ weight gain**).

### **Measurements**

The following measurements were recorded after calculation: average daily gain (ADG, g/d) was calculated as (**final BW-initial BW/days on feed**); intake was calculated(**feed offered – feed refused**); feed conversion ratio (FCR) was calculated by dividing **DMI/weight gain**; water consumption per unit feed intake (l/kg) was calculated as total water intake/dry matter intake (**TWI/DMI**); weight gain – the total weight an animal has gained for a given period of time(**ending weight – starting weight**); dressing percentage was calculated as (**hot carcass weight/live weight \* 100**).

## Statistical analysis

Data was analyzed using the General Linear Model procedures of MINITAB (2004). Significant differences among treatment means for dietary effect on growth parameters was analyzed using the least significant difference (LSD) method. Significance was declared at  $p < 0.05$ .

## Results

The chemical components of the commercial concentrate (ruminant mix) and mixed grass fed to growing lambs is presented in Table I. There was no difference ( $p > 0.05$ ) in dry matter (DM) (901 & 914 g/kg DM) and organic matter (OM) (886 & 891 g/kg DM) content between mixed grass samples and the commercial concentrate. Ash content was higher ( $p < 0.05$ ) in the mixed grass samples (104 g/kg DM) when compared to the commercial concentrate (79 g/kg DM).

**Table I Chemical composition of feeds used for feeding lambs at ECIAF (g/kg DM)**

Sample	DM g/kg	OM g/kg	Ash g/kg	CP g/kg	EE g/kg	CF g/kg	NFE g/kg
Mixed grass	901a	886a	104a	75.4 <sup>a</sup>	18.1 <sup>a</sup>	385a	418a
Ruminant mix	914a	891a	79b	135b	24.7 <sup>b</sup>	50.2 <sup>b</sup>	637b
SEM	0.18	0.10	0.09	0.22	0.32	1.55	3.09

DM: Dry matter, OM: Organic matter, CP: Crude protein, EE: Ether extract, CF: Crude fibre, NFE: Nitrogen

Free Extract

Calculated according to (AOAC 1995)

Crude protein (CP) levels were highest ( $p < 0.05$ ) in the commercial concentrate (135 g/kg DM) and lowest ( $p < 0.05$ ) in the mixed grass (75.4 g/kg DM). Similarly, ether extract (EE) was higher ( $p < 0.05$ ) in the commercial concentrate (24.7 g/kg DM) when compared to the mixed grass (18.1 g/kg DM). Crude fibre (CF) values were higher ( $p < 0.05$ ) in mixed grass samples (385 g/kg DM) when compared to the commercial concentrate (50.2 g/kg DM). In contrast, nitrogen free extract (NFE) was higher ( $p < 0.05$ ) in the commercial concentrate (637 g/kg DM) when compared to mixed grass (418 g/kg DM). The effect of different levels of concentrate supplementation on feed intake, water intake, average daily gain (ADG), feed conversion ratio (FCR) and weight gain of ewe lambs is presented in Table II. Feed intake was highest ( $p < 0.05$ ) among lambs supplemented at 3 % (5.85 kg) of their body weight. However, there was no significant difference ( $p > 0.05$ ) between the groups supplemented at 2 % (4.11 kg) and 2.5 % (4.72 kg) of their body weight. Similarly, there was no difference ( $p > 0.05$ ) in water intake between the groups supplemented at 2 % (12.9 L) and 2.5 % (12.9 L). However, water intake was highest ( $p < 0.05$ ) among animals supplemented at 3 % (15.3 L) of their body weight (Table II).

**Table II Effect of supplementation on the performance parameters of ewe lambs**

Treatment	Feed intake (kg)	Water intake (L)	ADG (kg)	FCR (kg)	Weight gain (kg)
2%	4.11 <sup>a</sup>	12.9 <sup>a</sup>	0.56 <sup>a</sup>	7.52 <sup>a</sup>	14.4 <sup>a</sup>
2.5%	4.72 <sup>a</sup>	12.9 <sup>a</sup>	0.56 <sup>a</sup>	8.50 <sup>b</sup>	16.3 <sup>ab</sup>
3%	5.85 <sup>b</sup>	15.3 <sup>b</sup>	0.70 <sup>b</sup>	8.28 <sup>b</sup>	20.6 <sup>c</sup>
SEM	0.189	1.044	0.021	0.210	1.662

<sup>abc</sup>Means in the same column within a parameter with different superscripts differ significantly;  $p < 0.05$ .

ADG:

Average daily gain, FCR: Feed conversion ratio, SEM: Standard error of mean

Average daily gain (ADG) was highest ( $p < 0.05$ ) among ewes supplemented at 3 % (0.70 kg) of their body weight. There was no difference ( $p > 0.05$ ) in average daily gains (ADG) between the groups supplemented at 2 % (0.56 kg) and 2.5 % (0.56 kg) respectively.

Feed conversion ratio (FCR) was significantly lower among animals supplemented at 2 % (7.52 kg) when compared to animals supplemented at 2.5 % (8.50 kg) and 3 % (8.28 kg) of their body weight respectively. Weight gain was significantly higher ( $p < 0.05$ ) among ewes supplemented at 3 % body weight in comparison to ewes supplemented at 2 % (14.4 kg) and 2.5 % (16.3 kg) respectively (Table II). The effect of concentrate supplementation on final weight, feed cost per pound of gain, dressing percentage and mortality of ewe lambs is illustrated in Table III. Final weight gained was significantly lower ( $p < 0.05$ ) among lambs supplemented at 2 % (34.8 kg) when compared to those supplemented at 2.5 % (46.4 kg) and 3 % (44.8 kg) of their body weight.

**Table III Effect of supplementation on weight, cost/gain and dressing percentage of ewe lambs**

Treatment	Final weight (kg)	Feed cost/lb of gain (\$TT)	Dressing percent (%)	Mortality (%)
2%	34.8 <sup>a</sup>	8.18 <sup>a</sup>	46 <sup>a</sup>	0
2.5%	46.4 <sup>b</sup>	9.52 <sup>a</sup>	45.7 <sup>a</sup>	0
3%	44.8 <sup>b</sup>	7.71 <sup>ab</sup>	42.9 <sup>b</sup>	0
SEM	1.662	1.133	0.524	-

<sup>abc</sup>Means in the same column within a parameter with different superscripts differ significantly;  $p < 0.05$ .

SEM: Standard error of mean



Ewes supplemented at 3 % of their body had a lower ( $p < 0.05$ ) feed cost per pound of gain (\$7.71) when compared to those supplemented at 2.5 % (\$ 9.52). Dressing percentage was also lowest ( $p < 0.05$ ) among ewes supplemented at 3 % of their body weight (42.9 %) when compared to ewes supplemented at 2 % (46 %) and 2.5 % (45.7 %) respectively. There was no difference ( $p > 0.05$ ) in mortality of lambs across the groups (Table III).

## **Discussion**

Grass forages in Trinidad and Tobago do not provide adequate protein to support the growing needs of lambs. This study emphasized the need to support local grass pastures with some level of concentrate. The crude protein (CP) levels of the mixed grass (CP 75.4 g/kg DM) as seen in Table I, would prove inadequate for fattening lambs as their CP requirement ranges from 130 to 160 g/kg DM (NRC, 2001). This mixed grass can be classified as a low quality forage. According to Obeidat *et al.* (2019), low quality forages are deficient in crude protein (less than 8% crude protein; CP), low in soluble sugars and starches, and are made up of natural pastures or crop residues. Supplementing the mixed grass with the National Feed Mills (NFM) Ruminant mix (CP 135 g/kg DM, Table I) have shown that the combination is more than capable of meeting the CP needs of these animals. The crude fibre (CF) content of the mixed grass is relatively high (385 g/kg DM, Tables I) which can lead to reduced intake due to gut fill. Additionally, high levels of fibre in the diet can result in decreased digestibility of dry matter thus limiting the supply of energy required for animal productivity. The crude fat (Ether extract) levels appear to be low (18-23 g/kg DM, Table I) for the mixed grass and the commercial concentrate. These levels may be adequate for rumen health as high levels of dietary fat (> 5 %) can negatively impact rumen microbes which can result in reduced degradability and dry matter intake owing to their inactivity.

Feed intake was highest ( $p < 0.05$ ) among lambs supplemented at 3 % (5.85 kg) of their body weight. This is in agreement with Hossain *et al.* (2003), Yagoub & Babiker (2008) and Sayed (2009) who reported that the level of energy significantly influenced daily feed intake of lambs, suggesting that nutrient digestibility increases with increased supplementation. Obeidat *et al.* (2019) concluded that lambs supplemented with crude protein (CP) will have greater feed intake and growth compared to the non-supplemented lambs grazing low-quality forages. Water intake was highest ( $p < 0.05$ ) among animals supplemented at 3 % (15.3 L) of their body weight (Table II). It is possible that as digestibility increases there is a greater demand for water to conduct metabolic processes. This study illustrated that average daily gain (ADG) was highest ( $p < 0.05$ ) among ewes supplemented at 3 % (0.70 kg) of their body weight. This supports the findings of (Asnakew, 2005) who indicated better gains in animals supplemented at high levels. Similarly, this finding agreed with that reported by Ebrahimiet *al.* (2007). They concluded that lambs fed on energy dense rations recorded higher significant average daily gain than that of lambs on medium and low energy diets. Also, average daily gain was highest in goat kids fed high-energy diet and lowest in goats fed low energy diets (Hossain *et al.*, 2003).

Increasing the levels of energy may facilitate the proliferation of more fermentable metabolizable energy (ME) for rumen microorganisms resulting in enhance microbial protein synthesis and increased amounts of protein available to the animal.

Final weight gained was significantly lower ( $p < 0.05$ ) among lambs supplemented at 2 % (34.8 kg) when compared to those supplemented at 2.5 % (46.4 kg) and 3 % (44.8 kg) of their body weight. This suggests that animals supplemented at higher levels are more efficient at utilizing feeds made available. Chegeniet *al.* (2019) and Manera *et al.* (2014) reported that the average daily gain (ADG) and the body weight (BW) gain of different sheep breeds were enhanced by those consuming higher levels of protein supplements.

Dressing percentage was also lowest ( $p < 0.05$ ) among ewes supplemented at 3 % of their body weight (42.9 %) when compared to ewes supplemented at 2 % (46 %) and 2.5 % (45.7 %) respectively. This does not support the findings of Payne and Wilson (1999) who reported that dressing percentage increases with increasing the proportions of concentrate in the ration. However, the dressing percentages of this study fell within the range of those reported by Macedo, Siqueira and Martins (1999). They concluded that the carcass yield of sheep may vary from 40 to 50%. The carcass yield obtained in this study was higher than those reported by Furusho-Garcia *et al.* (2010) (39.46%) in a study involving Santa Inês sheep slaughtered at an average weight of 38 kg. Ewes supplemented at 3 % of their body had a lower ( $p < 0.05$ ) feed cost per pound of gain (\$7.71) when compared to those supplemented at 2.5 % (\$ 9.52). The higher economic efficiency for lambs fed on high-energy diet compared with other supplemented groups could be attributed to increasing energy levels in lamb's diet. High energy diets are known to result in increased growth performance, adequate digestibility of nutrients and carcass yield of lambs (Sayed, 2009). Moreover, supplementing low-quality forage in ruminant diets with quality crude protein can reduce feeding expenses while maintaining or enhancing animal performance, voluntary feed intake and the digestibility of nutrients (Huston *et al.*, 1999 and Currier *et al.*, 2004).

## **Conclusions**

Generally, the present study demonstrated that supplementation of growing lambs with varied levels of concentrate on grass basal diets can improve feed intake, average daily gain, efficiency of feed conversion, weight gain and cost/lb of gain. Moreover, it was concluded that inclusion of concentrates at 3 % of lamb body weight resulted in better feed intake, daily gain, feed conversion efficiency and body weight of growing lambs compared to other rates of inclusion and could be recommended in practice. Further work should be conducted in an effort to establish an optimum inclusion rate.

## **Acknowledgements**

We extend sincere gratitude to the Programme Professor for allowing us access to the lambs and an experimental space at the ECIAF (UTT) campus Farm for execution of this study. We are indebted to the students who assisted with weighing, feeding and general husbandry practices. We are grateful that we were able to conduct our laboratory work at the Biosciences Agriculture and Food Technology (BAFT) laboratory of the University of Trinidad and Tobago. We thank the Laboratory Supervisor for facilitating us throughout the study.

**Author Contributions:** Andell Edwards - Conceptualization, Project Administration, Research design, Methodology, Data curation and writing of original draft. Amellia Taylor & Tunika Brewster – Daily husbandry, management and data collection. All authors have read and agreed to this draft.

**Conflicts of Interest:** The authors declare that there is no conflict of interest regarding this study.

## **References**

- Asnakew, A.(2005). Feedlot fattening performance and carcass characteristics of intact male Hararghe highland goats fed different levels of hay to concentrate ratios. M.Sc. Thesis, Alemaya University, Ethiopia. <https://core.ac.uk/download/pdf/132631371.pdf>
- AOAC.(1995).Official methods of analysis. 15th ed. Washington, DC, USA: Association of official Analytical Chemists.
- Chegini, R.Bonchenari, M.Farahani, A.H.Motlagh, M. Salem, A.Z.M. (2019). Effect of liquid protein feed on growth performance and ruminal metabolism of growing lambs fed low-quality forage and compared to conventional protein sources. J. Agric. Sci. 157: 272–280.<https://doi.org/10.1017/S0021859619000595>.
- Currier, T.A.Bohnert, D.W. Falckv, S.J. Bartle, S.J.(2004).Daily and alternate day supplementation of urea or biuret to ruminants consuming low-quality forage: II. Effects on cow performance and efficiency of nitrogen use in wethers. J. Anim. Sci. 82,1508–1517.<https://doi.org/10.2527/2004.8251508x>.
- Ebrahimi, R. et.al. (2007). Effect of energy and protein levels on feedlot performance and carcass characteristics of Mehraban ram lambs. Pak. J. Biol.Sci.15(10), 1679-1684.<https://doi.org/10.3923/pjbs.2007.1679.1684>
- Furusho-Garcia I.F. et al. (2010).Performance and carcass characteristics of Santa Inês pure lambs and crosses with Dorper and Texel at different management systems. Revista Brasileira de Zootecnia. 39(6), 1313- 1321.<https://doi.org/10.1590/S1516-35982010000600021>
- Hossain M.E.Shahjalal, M. Khan, M.J.Hasanat, M.S. (2003).Effect of dietary energy supplementation on feed intake, growth and reproductive performance of goats under grazing condition. Pak. J. of Nutr. 2(3), 159163.<https://dx.doi.org/10.3923/pjn.2003.159.163>
- Huston, J.E.Engdahl, B.S. Bales K.W. (1999). Supplemental feeding interval for adult ewes. Sheep Goat Res. Jour. 15, 87–93.<http://dx.doi.org/10.3390/ani10010051>
- Khanal, R.C.Gurung D.B.Kadariya R.K. (1999). Effect of feeding urea treated rice and wheat straw on intake and milk yield of lactating buffaloes under farmers conditions. Asian-Australasian Journ of Anim Sci. 12(8),1200-1204. <https://doi.org/10.5713/ajas.1999.1200>



- Kirton, A.H. Carter, A.H. Clarke, J.N. et al. (1995b). A comparison between 15 ram breeds for export lamb production. 1. Liveweights, body components, carcass measurements, and composition. N.Z. J. Agric. Res. 38, 347. <https://doi.org/10.1080/00288233.1996.9513193>
- Macedo, F.A.F.Siqueira, E.R. Martins, E.N. (1999).Desempenho de cordeiros corriedale puros e mestiços, terminados em pastagem e em confinamento. Arquivo Brasileiro de Medicina Veterinária e Zootecnia. 51(6), 583-587. <https://doi.org/10.1590/S0102-09351999000600014>.
- Manera, D.B.Voltolini, T.V. Yamamoto, S.M. Araujo, G.G. Souza R.A. (2014). Productive performance lambs on grazing supplemented with concentrates containing fruit processing by-products. Semin. Cienc. Agrar. 35, 1013–1022. <http://dx.doi.org/10.3390/ani10010051>
- McDonald, P. Edwards, R.A. Greenhalgh J.F.D. Morgan, C.A. (2002). Animal Nutrition, sixth ed. Pearson Educational Limited, Harlow, UK.
- MuluMoges (2005). Effect of feeding different levels of breweries' dried grain on live weight and carcass characteristics of Wogera sheep fed on hay basal diet. MSc Thesis presented to school of graduate studies, Haramaya University, 56p. <http://www.lrrd.org/lrrd23/4/take23072.htm>
- National Research Council (2001). Nutrient requirements of sheep. 6. ed. Washington: National Academy Press. 99p.
- Nega, A. and Melaku, S. (2009). Feed intake, digestibility and body weight change in Farta sheep fed hay supplemented with rice bran and/or noug seed (*Guizotia abyssinica*) meal. Trop Anim Health and Prod. 41(4), 507-515. <https://doi.org/10.1007/s11250-008-9138-1>
- Obeidat, B.S. Subih, H.S. and Ata, M. (2019). Protein Supplementation Improves Performance of Lambs Fed Low Quality Forage. Animals (Basel). 10(1), 51. doi: 10.3390/ani10010051. PMID: 31881754; PMCID: PMC7022730.
- Payne, W.J.A. and Wilson, R.T. (1999). An introduction to animal husbandry in the tropics (No. Edn 5). Blackwell Science.
- Rege, J.E.O. (1999). "Economic valuation of farm animal genetic resources." In Proceedings of an FAO/ILRI Workshop, Roma, pp. 25-27. 10.3920/978-90-8686-592-5
- Sandrose, D. (1993). Comparative study on growth, carcass and wool traits in Menz sheep and their crosses with the Awasia and Corriedale breeds under supplemented grazing. M.Sc. Thesis, Alemaya University, Ethiopia.

- Sayed, A.B.N. (2009). Effect of different dietary energy levels on the performance and nutrient digestibility of lambs. *Vet. World.* 2, 418-420. <https://doi=10.1.1.302.5831&rep=rep1&type=pdf>
- Simret, B.(2005). Supplementation of graded levels of peanut cake and wheat bran mixture on nutrient utilization and carcass parameters of Somali goats. M.Sc. Thesis Presented to the School of Graduate Studies of Alemaya University of Agriculture, Alemaya, Ethiopia. Pp. 60.
- Van Soest, P.J. Robertson, J.B. and Lewis, B.A. (1991). Methods for dietary fiber, neutral detergent fiber, and nonstarch polysaccharides in relation to animal nutrition. *J. Dairy Sci.* 74, 3583-3597. [https://doi.org/10.3168/jds.S0022-0302\(91\)78551-2](https://doi.org/10.3168/jds.S0022-0302(91)78551-2)
- Van Soest, P.J. (1994). *Nutritional ecology of the ruminant.* Cornell university press.
- Yagoub, Y.M. and Babiker, S.A. (2008). Effect of dietary energy level on growth and carcass characteristics of female goats in Sudan. *J. Anim. Sci.* 80, 100-115. <http://www.lrrd.org/lrrd20/12/yago20202.htm>
- Zemichael, G. (2007). Supplementation of sesame seed (*Sesame indicum*) cake, wheat bran and their mixtures on feed intake, digestibility, live weight changes and carcass characteristics of Tigray sheep fed a basal diet of tef straw. MSc Thesis, Haramaya University, Ethiopia.